

Theraderm 2500™

Pre & Post Sanitizing Teat Dip

Theraderm 2500 is a .25% (2500 ppm) iodine teat dip formulated to be effective for use in pre- and post-dipping applications, where a lower titratable iodine product is desired.

Features and Benefits

- **Lower Iodine Level**

Excellent for use on dairy farms that pre- and post-dip 2 or 3 times per day, and for cows with a low tolerance to iodine.

- **Exclusive Triple Emollient System**

Contains lanolin, glycerin and a special therapeutic ingredient to promote a natural moisture balance that keeps teats soft and pliable.

- **Contains Surfactant**

Provides excellent wetting qualities to help coat the teat and clean residual soils from teats when used as a pre-dip.

- **Aids in Decreasing Mastitis**

Thoroughly tested iodine formula provides quick, broad-spectrum kill of bacteria, reducing the chance of new intramammary infections.

- **Controlled pH**

Provides kill of bacteria without danger of irritation to cows' teats or operators' hands.

- **Ready to Use**

No mixing is required.

- **Freeze/Thaw Stability**

If frozen, this product can be thawed completely, mixed to its original state and used with confidence.

- **100% Customer Satisfaction**

All GEA Farm Technologies hygiene products have a 100% Customer Satisfaction Guarantee.



GEA Farm Services | WestfaliaSurge



Theraderm 2500™

Pre & Post Sanitizing Teat Dip

Directions for Use

Use at full strength. Do not dilute.

Indications: This topical liquid product, when used undiluted as a pre- and post-milking teat dip, effectively aids in reducing the spread of mastitis.

Pre Dipping: Before milking, dip each teat with Theraderm 2500. After 15 to 30 seconds, dry each teat thoroughly with a single service towel. If the udder and teats are heavily soiled, wash with a sanitizing solution and dry before application of Theraderm 2500.


Post Dipping: Immediately after milking, dip each teat with Theraderm 2500. Allow teats to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before the cow leaves the parlor to prevent freezing. At the end of lactation, apply this product daily for one week after the last milking. In addition, begin application of this product about one week prior to calving.

If a common teat dip cup is used for application, a fresh solution should always be used at each milking. The teat dip cup should be emptied, cleaned and rinsed with potable water after each milking session or when cup becomes contaminated during milking. Do not pour remaining solution from dip cup back into original container.

Research

Theraderm 2500 has been tested extensively in germicidal activity tests, skin irritation studies and university studies (Teat Dip Protocols) consistent with those recommended by the NMC. The results of all studies proved Theraderm 2500 to be an excellent mastitis prevention tool.

Complete research data available upon request.



GEA Farm Technologies

GEA Farm Technologies, Inc.

1880 Country Farm Drive, Naperville, IL 60563
Toll Free: 1.877.WS.DAIRY or 1.877.973.2479
www.gea.com | www.gea-farmtechnologies.com

Product Properties

Appearance: Dark red-brown

Odor: Iodine

pH: 5.3

Specific Gravity at 77°F (25°C): 1.03 (water = 1)

Wetting Ability: Excellent

Ordering Information

U.S. (only)

SKU	Description
7751-0041-511	Theraderm 2500 - 5 gal. pail (18.9 liters)
7751-0041-510	Theraderm 2500 - 15 gal. drum (56.7 liters)
7751-0041-508	Theraderm 2500 - 55 gal. drum (208 liters)
7751-1000-434	Theraderm - Literature (pkg of 20)

7751-1000-434 (pkg of 20)
(rev. 07-13)