

Theratec[®]

Pre & Post Sanitizing Teat Dip

Theratec is a ready-to-use 0.5% (5000 ppm) iodine teat dip, formulated to be safe and effective for use in pre- and post-dipping applications (spraying or dipping).

Features and Benefits

- **Finishes What It Starts**

The 0.5% formulation of Theratec provides the right iodine level for both pre- and post-dipping applications.

- **Aids in Decreasing Mastitis Cases**

Thoroughly tested iodine formula provides a quick, broad-spectrum kill of bacteria, reducing the chance of new intramammary infections.

- **Exclusive Triple Emollient System**

Contains lanolin, glycerin and a special therapeutic ingredient to promote a natural moisture balance that keeps teats soft and pliable.

- **Contains Surfactants**

Provides excellent wetting qualities to help coat the teat and clean residual soils from teats when used as a pre-dip.

- **Controlled pH**

Provides kill of bacteria without danger of irritation to cows' teats or operators' hands.

- **Ready to Use**

No mixing is required.

- **Freeze/Thaw Stability**

If frozen, this product can be thawed completely, mixed to its original state and used with confidence.

- **100% Customer Satisfaction**

All GEA Farm Technologies hygiene products have a 100% Customer Satisfaction Guarantee.



GEA Farm Services | WestfaliaSurge



Theratec®

Pre & Post Sanitizing Teat Dip

Directions for Use

Use at full strength. Do not dilute.

Indications: This topical liquid product, when used undiluted as a pre- and post-milking teat dip, effectively aids in reducing the spread of mastitis.

Pre-Dipping: Before milking, dip each teat with Theratec. After 15 to 30 seconds, dry each teat thoroughly with a single service towel. If the udder and teats are heavily soiled, wash with a sanitizing solution and dry before application of Theratec.

Post-Dipping: Immediately after milking, dip each teat with Theratec. Allow teats to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teats before the cow leaves the parlor to prevent freezing. At the end of lactation, apply this product daily for one week after the last milking. In addition, begin application of this product about one week prior to calving.

If a common teat dip cup is used for application, a fresh solution should always be used at each milking. The teat dip cup should be emptied, cleaned and rinsed with potable water after each milking session or when cup becomes contaminated during milking. Do not pour remaining solution from dip cup back into original container.

Research

Theratec has been tested extensively in germicidal activity tests, skin irritation studies and university studies (Teat Dip Protocols) consistent with those recommended by the NMC. The results of all studies proved Theratec to be an excellent mastitis prevention tool.

Complete research data available upon request.

Product Properties

- Appearance:** Dark red/brown liquid
- Odor:** Mild iodine
- pH:** 5.3 (as is)
- Specific Gravity at 77°F (25°C):** 1.02 (water = 1)
- Wetting Ability:** Excellent


Ordering Information

U.S. (only)


SKU	Description
7751-0240-485	Theratec - Case (4x1 gal.) (4x3.78 liters)
7751-0040-487	Theratec - 5 gal. pail (18.9 liters)
7751-0040-488	Theratec - 15 gal. drum (56.7 liters)
7751-0040-489	Theratec - 30 gal. drum (113.5 liters)
7751-0040-649	Theratec - 55 gal. drum (208 liters)
7751-1000-431	Theratec - Literature (pkg of 20)




Experimental Challenge: Cornell University



Experimental Challenge: Louisiana State University



Natural Exposure Testing: Louisiana State University



GEA Farm Technologies

GEA Farm Technologies, Inc.

1880 Country Farm Drive, Naperville, IL 60563
Toll Free: 1.877.WS.DAIRY or 1.877.973.2479
www.gea.com | www.gea-farmtechnologies.com

7751-1000-431 (pkg of 20)
(rev. 07-13)